

# Immunization Site Map

Suggested sites for  
infant immunizations:



RD: \_\_\_\_\_

LD: \_\_\_\_\_

RT: \_\_\_\_\_

LT: \_\_\_\_\_

RT: \_\_\_\_\_

LT: \_\_\_\_\_

RD= Right deltoid (IM) or subcutaneous  
tissue on upper arm (SC).

RT= Right vastus lateralis (IM) or subcuta-  
neous tissue on thigh (SC).

LD= Left deltoid (IM) or subcutaneous  
tissue on upper arm (SC).

LT= Left vastus lateralis (IM) or subcu-  
taneous tissue on thigh (SC).



California Department of Health Services • Immunization Branch • 2151 Berkeley Way • Berkeley, CA 94704



IMM-718 (5/01)

# Immunization Site Map

Suggested sites for toddler immunizations:



RD: \_\_\_\_\_

RT: \_\_\_\_\_

RT: \_\_\_\_\_

LD: \_\_\_\_\_

LT: \_\_\_\_\_

LT: \_\_\_\_\_


RD= Right deltoid (IM) or subcutaneous tissue on upper arm (SC).  
RT= Right vastus lateralis (IM) or subcutaneous tissue on thigh (SC).

LD= Left deltoid (IM) or subcutaneous tissue on upper arm (SC).  
LT= Left vastus lateralis (IM) or subcutaneous tissue on thigh (SC).




# Giving All The Doses

## One way to give 5 doses at one visit



## One way to give 7 doses (if needed) at one visit



The deltoid muscle is an option for IM injections in children 18 months and older with adequate muscle mass.