

Injection Safety

Test Your Knowledge-Template

This document can be used as part of the overall competency program when conducting competency evaluation and/or validation of healthcare personnel (HCP) for injection safety.

Injection Safety Assessment	True	False
HCP who prepares and/or administers parenteral medications are required to receive training on safe injection practices upon hire and annually.	<input type="checkbox"/> True	<input type="checkbox"/> False
Changing the needle makes it safe to re-use a syringe.	<input type="checkbox"/> True	<input type="checkbox"/> False
Syringes can be re-used as long as the injection is administered through IV tubing.	<input type="checkbox"/> True	<input type="checkbox"/> False
Multi-dose vials should be discarded after 28 days from date of opening/puncture, unless manufacturer specifies otherwise.	<input type="checkbox"/> True	<input type="checkbox"/> False
It is acceptable to use leftover medicine from a single-dose or single-use vial for more than one patient.	<input type="checkbox"/> True	<input type="checkbox"/> False
Injections are to be prepared using aseptic technique in a clean area free from contamination or contact with blood, body fluids or contaminated equipment.	<input type="checkbox"/> True	<input type="checkbox"/> False
Multi-dose vials should be stored inside the immediate patient care area and under manufacturer storage recommendations.	<input type="checkbox"/> True	<input type="checkbox"/> False
Sharps do not have to be in a puncture-resistant sharps container.	<input type="checkbox"/> True	<input type="checkbox"/> False