

Animal Bite/Potential Rabies Exposure Surveillance and Investigation Protocol

Table of Contents

I.	ABOUT THE DISEASE	2
A.	Clinical Presentation	2
B.	Etiologic Agent	2
C.	Reservoir	2
D.	Incubation Period	2
E.	Mode of Transmission	3
F.	Period of Communicability	3
II.	DISEASE CONTROL AND PREVENTION	3
A.	Disease Control Objectives	3
B.	Disease Prevention Objectives	3
C.	Disease Prevention and Control Intervention	3
D.	Treatment	7
III.	DISEASE INVESTIGATION	7
A.	Case Definition and Case Classification	7
B.	Reporting Timeframe to Public Health	8
C.	Outbreak Recognition	8
D.	Healthcare Provider Responsibilities	8
E.	Laboratory Responsibilities	8
F.	Local Health Responsibilities	9
G.	State Health Responsibilities	10
H.	Occupational Health	10
IV.	DISEASE SURVEILLANCE	10
A.	Public Health Significance	10
B.	Disease Surveillance Objectives	11
C.	Surveillance Indicators	11
V.	REFERENCES	13

Office of Epidemiology and Prevention Services

Division of Communicable Disease Epidemiology

350 Capitol Street Room 125, Charleston, WV 25301-3715

Phone: (304) 558-5358 ext. 2 Fax: (304) 558-6335 www.oeps.wv.gov

Animal Bite/Potential Rabies Exposure Surveillance and Investigation Protocol

I. ABOUT THE DISEASE

Rabies is a preventable viral disease most often transmitted through saliva from the bite of a rabid mammal animal.¹ Raccoon strain rabies is endemic in West Virginia, making animal bite reporting the first step in preventing human disease.²

A. Clinical Presentation

Clinical signs of rabies start out like the flu including general weakness or discomfort, fever, or headache. There may be discomfort around the bite area as well as cerebral dysfunction, anxiety, confusion, and agitation. As the disease progresses, the person may experience delirium, abnormal behavior, hallucinations, hydrophobia, and insomnia. The acute period lasts 2 to 10 days and once clinical signs appear the disease is almost always fatal.

B. Etiologic Agent

The rabies virus belongs to the order Mononegavirales, viruses with nonsegmented, negative stranded RNA genomes. Within this group, viruses with a distinct "bullet" shape are classified in the *Rhabdoviridae* family, which includes at least three genera of animal viruses, *Lyssavirus*, *Ephemerovirus*, and *Vesiculovirus*. The genus *Lyssavirus* includes the rabies virus.^{12,13} Several other pathogenic organisms may also be present in the saliva of an exposing animal, including but not limited to *Pasteurella multocida*, *Eikenella corrodens*, *Capnocytophaga spp*, *Neisseria weaveri*, *Weeksella zoohelcum*, *Neisseria canis*, and *Staphylococcus intermedius*.^{14,15}

C. Reservoir

Although all species of mammals are susceptible to rabies virus infection, only a few species are important as reservoirs for the disease.¹³ In the United States, several distinct rabies virus variants have been identified in terrestrial mammals, including raccoons, skunks, foxes, and coyotes. In addition to these terrestrial reservoirs, several species of insectivorous bats are also reservoirs for rabies. Raccoons serve as the primary reservoir of rabies in West Virginia.^{2,16}

D. Incubation Period

The incubation period for rabies usually ranges from 1 to 3 months after exposure but can range from a few days to several years. The type and intensity of exposure and location of the wound partly influences the incubation period.¹³

Office of Epidemiology and Prevention Services

Division of Communicable Disease Epidemiology

350 Capitol Street Room 125, Charleston, WV 25301-3715

Phone: (304) 558-5358 ext. 2 Fax: (304) 558-6335 www.oeps.wv.gov

Animal Bite/Potential Rabies Exposure Surveillance and Investigation Protocol

E. Mode of Transmission

Transmission of rabies virus usually begins when infected saliva of a host is passed to an uninfected animal. Various routes of transmission have been documented and include contamination of mucous membranes (i.e., eyes, nose, and mouth), aerosol transmission, and corneal or other tissue transplants. The most common mode of rabies virus transmission is through virus-containing saliva contaminating the bite from an infected host. ¹³

F. Period of Communicability

Rabies virus may be excreted in the saliva of infected dogs, cats, and ferrets during illness and/or 3 - 7 days prior to illness or death; these are the only domestic animals with defined, reliable periods of communicability. ¹³ The incubation period and number of days the rabies virus is shed in the saliva, prior to onset of clinical signs, are unknown for wild and exotic animals. Infected animals can transmit the rabies virus for an indeterminate number of days before clinical signs become apparent, as well as while clinically symptomatic. ^{1, 12, 13} **Therefore, it is never appropriate to confine and observe wild or exotic animals that are involved in a human bite incident.**

II. DISEASE CONTROL AND PREVENTION

A. Disease Control Objectives

Prevent human rabies through rapid identification of cases potentially exposed to the rabies virus so appropriate treatment or post exposure prophylaxis (PEP) can be quickly administered.

B. Disease Prevention Objectives

To prevent rabies transmission through:

1. Education of the public about animal bite prevention and vaccination of dogs, cats, and ferrets.
2. Education of the public on animal bite reporting and seeking medical attention.
3. Education of physicians and other public health partners on appropriate management of animal bite cases.

C. Disease Prevention and Control Intervention

1. Educate the public about pre-exposure vaccination if traveling to a country with endemic animal rabies.
2. Educate the public on vaccinating pets and keeping vaccinations up to date.

Office of Epidemiology and Prevention Services

Division of Communicable Disease Epidemiology

350 Capitol Street Room 125, Charleston, WV 25301-3715

Phone: (304) 558-5358 ext. 2 Fax: (304) 558-6335 www.oeps.wv.gov

Animal Bite/Potential Rabies Exposure Surveillance and Investigation Protocol

3. Educate the public on avoiding stray or wild animals. Wild/domestic crossbreeds (ex. dog/wolf) should not be kept as pets.
4. Educate the public that if bitten, scratched, or unsure, talk to a healthcare provider about the need for postexposure prophylaxis. Prompt wound care and administration of rabies immune globulin (RIG) and vaccine are highly effective in preventing human rabies following exposure.
5. Veterinarians and veterinary support staff, who are frequently at risk for rabies exposures, should obtain pre-exposure rabies vaccinations, followed by periodic titer checks and rabies vaccine boosters. The ACIP recommends titers on a schedule of every two years to assess protective immunity, with a single-injection booster vaccination recommended if the titer level is below a 1:5 serum dilution (0.1-0.2 IU/mL).
6. Educate clinical personnel to use standard precautions when providing care to persons suspected of having clinical rabies, including wearing gowns, goggles, masks, and gloves, particularly during procedures that might result in splashes or sprays from body fluids.
7. For animal bites, wound care and cleaning is important to prevent infection and, depending on the status and species of the animal in conjunction with the type of exposure, postexposure prophylaxis (PEP) may be warranted.^{1, 3, 4,13}

For guidance on PEP:

1. Rabies Risk Exposure for Human Exposure to Animals Algorithm for PEP, available [here](#).
2. ACIP recommendations on PEP, available [here](#).¹⁷
3. Summary of rabies postexposure prophylaxis (PEP) recommendations from CDC [here](#)¹ and below. For unvaccinated persons, the combination of RIG and vaccine is recommended for both bite and non-bite exposures, regardless of the time interval between exposure and initiation of PEP. If PEP has been initiated and appropriate laboratory diagnostic testing (direct fluorescent antibody test) indicates the animal that caused the exposure was not rabid, PEP may be discontinued.

Office of Epidemiology and Prevention Services
Division of Communicable Disease Epidemiology

350 Capitol Street Room 125, Charleston, WV 25301-3715

Phone: (304) 558-5358 ext. 2 Fax: (304) 558-6335 www.oeps.wv.gov

Animal Bite/Potential Rabies Exposure Surveillance and Investigation Protocol

Postexposure Prophylaxis for Non-immunized Individuals

Treatment	Regimen
Wound cleansing	All postexposure prophylaxis should begin with immediate thorough cleansing of all wounds with soap and water. If available, a virucidal agent such as povidine-iodine solution should be used to irrigate the wounds.
RIG	If possible, the full dose should be infiltrated around any wound(s) and any remaining volume should be administered IM at an anatomical site distant from vaccine administration. Also, RIG should not be administered in the same syringe as vaccine. Because RIG might partially suppress active production of antibody, no more than the recommended dose should be given.
Vaccine	HDCV or PCECV 1.0 mL, IM (deltoid area), one each on days 0 , 3, 7, and 14.

** A 5th dose on day 28 may be recommended for immunocompromised persons.*

Postexposure Prophylaxis for Previously Immunized Individuals

Treatment	Regimen
Wound cleansing	All postexposure prophylaxis should begin with immediate thorough cleansing of all wounds with soap and water. If available, a virucidal agent such as povidine-iodine solution should be used to irrigate the wounds.
RIG	RIG should not be administered.
Vaccine	HDCV or PCECV 1.0 mL, IM (deltoid area), one each on days 0 and 3.

If exposed to rabies, previously vaccinated persons should receive two IM doses (1.0 mL each) of vaccine, one immediately and one three days later. Previously vaccinated persons are those who have received one of the recommended preexposure or postexposure regimens of HDCV, RVA, or PCECV, or those who received another vaccine and had a documented rabies antibody titer. RIG is unnecessary and should not be administered to these persons because an anamnestic response will follow the administration of a booster regardless of the pre-booster antibody titer.

- Every attempt should be made to adhere to the recommended vaccination guidelines. If a dose is missed by a few days, give the next dose as soon as possible and give the subsequent doses in the appropriate # of days later as typically indicated.¹
 - Dose 1= day 0
 - Dose 2= 3 days later
 - Dose 3= 4 days later
 - Dose 4= 7 days later

Office of Epidemiology and Prevention Services
Division of Communicable Disease Epidemiology
 350 Capitol Street Room 125, Charleston, WV 25301-3715
 Phone: (304) 558-5358 ext. 2 Fax: (304) 558-6335 www.oeps.wv.gov

Animal Bite/Potential Rabies Exposure Surveillance and Investigation Protocol

- a) When substantial deviations from the schedule occur, immune status should be assessed by performing serologic testing 7-14 days after administration of the final dose in the series.¹
 - b) If rabies immunoglobulin (HRIG) is not available on the first visit, it can be given up to, and including, day 7 from the first vaccine dose. Once 7 days have passed the patient's antibody response to the vaccine occurs and can be altered if HRIG is given.¹
 - c) No routine testing of healthy patients completing PEP is necessary to document seroconversion.
 - d) For more information on delays/interruptions in PEP schedule see [here](#).
5. Information from CDC on rabies vaccine for uninsured/underinsured (indigent) population is available [here](#).
 6. Pregnancy is not considered a contraindication to PEP. Certain studies have indicated no increase in abortion, premature births, or fetal abnormalities associated with rabies vaccination. If the risk of rabies exposure is substantial, PEP might be indicated during pregnancy. Rabies exposure or diagnosis of rabies in the mother should not be regarded as reasons to terminate the pregnancy.¹
 7. For patients taking corticosteroids, other immunosuppressive agents, antimalarials, and those who are immunosuppressed, rabies PEP should be administered using a 5-dose vaccine regimen (i.e., 1 dose of vaccine on days 0, 3, 7, 14 and 28), with the understanding that immune response might be inadequate. Immunosuppressive agents should not be administered during rabies PEP unless essential for the treatment of other conditions. If possible, immunosuppressed patients should postpone rabies preexposure prophylaxis until the immunocompromising condition is resolved. When postponement is not possible, immunosuppressed persons who are at risk for rabies should have their virus neutralizing antibody responses checked after completing the preexposure series. Postvaccination rabies virus-neutralizing antibody values might be less than adequate among immunosuppressed persons with HIV or other infections. When rabies pre- or post-exposure prophylaxis is administered to an immunosuppressed person, one or more serum samples should be tested for rabies virus-neutralizing antibody by the RFFIT to ensure that an acceptable antibody response has developed after completing the series. If no acceptable antibody response is detected after the final dose in the pre- or postexposure prophylaxis series, the patient should be managed in consultation with their physician and appropriate public health officials.

Office of Epidemiology and Prevention Services

Division of Communicable Disease Epidemiology

350 Capitol Street Room 125, Charleston, WV 25301-3715

Phone: (304) 558-5358 ext. 2 Fax: (304) 558-6335 www.oeps.wv.gov

Animal Bite/Potential Rabies Exposure Surveillance and Investigation Protocol

Additional information about rabies surveillance management of suspect rabid animals following human and animal exposures:

https://oeps.wv.gov/rabies/documents/lhd/RABIES_GUIDE.pdf.

D. Treatment

No proven medical treatment exists for symptomatic human rabies and disease almost invariably results in death.

III. DISEASE INVESTIGATION

A. Case Definition and Case Classification

Human Exposure

A bite or scratch from a vector species or the introduction of saliva or central nervous system (CNS) tissue from a vector species into an open, fresh wound or mucous membrane (eye, mouth, or nose) of a human being.

Vector Species

Species include bats or terrestrial mammals, especially carnivores. Wild species known to be reservoirs of rabies include, but are not limited to, raccoons, skunks, foxes, coyotes, bobcats, wolves, or any hybrids between these wild species and domestic dogs and cats. Domestic species include, but are not limited to, dogs, cats, and ferrets.

Case Classifications

Confirmed: Human exposure from a vector species as defined above.

Other Classification Criteria

Touching or handling a potentially rabid animal or another animal or inanimate object that had contact with a rabid animal does not constitute an exposure unless wet saliva or CNS material from the rabid animal was introduced into a fresh, open wound or had contact with a mucous membrane of a human being.¹⁵

Bats have small teeth which may leave marks that are not easily seen; therefore, any contact with a bat in which a bite cannot be ruled out is considered a potential exposure to rabies. A person sleeping in a room with a bat or finding a bat in the room with an

Office of Epidemiology and Prevention Services

Division of Communicable Disease Epidemiology

350 Capitol Street Room 125, Charleston, WV 25301-3715

Phone: (304) 558-5358 ext. 2 Fax: (304) 558-6335 www.oeps.wv.gov

Animal Bite/Potential Rabies Exposure

Surveillance and Investigation Protocol

unattended child, person with mental disability, or intoxicated person, are examples of possible exposures.

Laboratory Diagnosis

The Office of Laboratory Services (OLS) is responsible for screening animal brain tissue for the presence of the rabies virus. This is the only facility in the state of West Virginia that can screen animal brain tissue for the presence of the rabies virus following human exposure. A direct fluorescent antibody (DFA) staining technique is used. The standard protocol for accepting suspect rabid animals includes animals involved with biting or scratching humans, animals involved with biting or scratching domestic animals or livestock, occasional environmental 'spot check' of areas (surveillance), and unusual situations involving the suspect animals such as atypical behavior.

Information for submitting specimens (including rabies specimen submission form) can be found at the OLS website: <https://dhhr.wv.gov/ols/labs/Pages/Rabies.aspx>.

B. Reporting Timeframe to Public Health

Animal bites and other potential rabies exposures must be reported to the local health department within 24 hours.

C. Outbreak Recognition

Outbreaks of animal rabies may be recognized as an increase in the number of cases over or above baseline incidence.

D. Healthcare Provider Responsibilities

1. Report any animal bite/exposure to the local health department of the victim's residence within **24 hours**.
2. Provide victim and exposing animal information using the Animal Encounter Report Form. This form can be found [here](#).
3. If necessary, consult with the local or state public health officers regarding the need for post exposure prophylaxis (PEP). Additional guidance is available in the Morbidity and Mortality Weekly Report (MMWR) (Volume 59, No. RR-2) linked [here](#).

E. Laboratory Responsibilities

1. All requests for animal rabies testing should be arranged through the local health department and the West Virginia Department of Health and Human Resources'

Office of Epidemiology and Prevention Services

Division of Communicable Disease Epidemiology

350 Capitol Street Room 125, Charleston, WV 25301-3715

Phone: (304) 558-5358 ext. 2 Fax: (304) 558-6335 www.oeps.wv.gov

Animal Bite/Potential Rabies Exposure Surveillance and Investigation Protocol

(DHHR) Office of Laboratory Services at (304) 558-3530, ext. 20135. Requests for human rabies testing are arranged through the Office of Epidemiology and Prevention Services (OEPS) at (304) 558-5358, ext. 2.

2. Report positive results to the Office Epidemiology and Prevention Services.

F. Local Health Responsibilities

1. Be familiar with *Animal Bite/Potential Surveillance and Investigation Protocol*.
2. Provide education on reporting requirements to local healthcare providers and other partners. Partners may include emergency room personnel, urgent care personnel, local physician offices, hospital infection control personnel, animal control officers and sheriff's department personnel. Animal bites are reportable to the local health department within 24 hours.
3. The [Animal Encounter Report Form](#) should be completed by healthcare providers and sent to the local health department. The local health department should enter bite encounters into WVEDSS (ensure data is complete and correct, see section IV for required surveillance indicators). Attach the completed form to the supplemental info tab in the WVEDSS investigation.
4. Upon receiving an animal encounter form, the local health department should determine circumstances of the bite encounter. The local health department should determine whether the animal is wild, stray, or owned; if owned the local health department should attempt to identify the owner.
5. The local health department should identify the confinement status of the animal, where needed. See [West Virginia Code](#). Confinement guidelines can be found in DC-4 available [here](#). After the confinement period is over, the LHD should follow up on the animal and document the outcome in WVEDSS. If the animal has been killed or has died, and the animal brain is in good condition and available for testing, the LHD should arrange rabies testing for the animal at the Office of Laboratory Services.
6. Contact the bite victim. The LHD will follow up with the victim after the confinement period or after laboratory results are known and document findings in WVEDSS, including rabies post-exposure prophylaxis and management.
 - i. Discuss exposure management with victim. See [Rabies Risk Assessment for Human Exposure to Animals](#). Provide recommendations for [Post Exposure Prophylaxis \(PEP\)](#). For assistance contact the OEPS epidemiologist on-call at (304) 558-5358, ext. 2. An epidemiologist is available 24/7/365.

Office of Epidemiology and Prevention Services

Division of Communicable Disease Epidemiology

350 Capitol Street Room 125, Charleston, WV 25301-3715

Phone: (304) 558-5358 ext. 2 Fax: (304) 558-6335 www.oeps.wv.gov

Animal Bite/Potential Rabies Exposure Surveillance and Investigation Protocol

- ii. To be considered lost to follow-up (LTFU), there must have been three attempts to call the victim, and a letter must have been sent with no reply.

G. State Health Responsibilities

1. Collect and review animal bite encounter reports for data completeness and quality.
2. Provide training on animal bite and other rabies exposure management to local health departments and other community health partners.
3. Develop evidence-based education related to animal bite and rabies exposure prevention for dissemination to the public.
4. Provide consultation to local health departments and community health partners on PEP recommendations and rabies exposure management.
5. In the event of a Herpes B Virus exposure, facilitate and arrange testing with the University of Georgia Herpes B Virus Resource Center in conjunction with the Office of Laboratory Services.

H. Occupational Health

1. Ensure all staff are trained on basic infection control principles and use universal precautions when handling animals for decapitation.
2. Ensure all staff have up-to-date vaccinations for other communicable diseases.
3. For staff performing decapitations, ensure that staff are currently immunized against rabies.
 - a) For proper personal protective equipment (PPE), see the DC-4 Appendix B.
4. For any staff who may be at risk of an animal exposure in the course of their duties, they should consider rabies pre-exposure prophylaxis vaccination.

IV. DISEASE SURVEILLANCE

A. Public Health Significance

One of every two Americans will be bitten by an animal at some point.³ Mammalian bites account for approximately 1% of all visits to emergency rooms, resulting in about 2 million bite wounds costing \$30 million dollars yearly.⁴ Annual mortality rate from dog bites is reported as 6.1 per 100 million population, based on a yearly average of 19 reported deaths from dog bites per year in the United States.^{5, 6}

Office of Epidemiology and Prevention Services

Division of Communicable Disease Epidemiology

350 Capitol Street Room 125, Charleston, WV 25301-3715

Phone: (304) 558-5358 ext. 2 Fax: (304) 558-6335 www.oeps.wv.gov

Animal Bite/Potential Rabies Exposure Surveillance and Investigation Protocol

Children are largely the victims of animal bites, particularly in cases involving serious injury.^{7,8}

Although 90% of animal bites are perpetrated by dogs and cats, most human rabies cases are caused by bat exposures.⁹ From 2009 – 2019, a total of 25 human cases were reported (with a 92.0% case-fatality rate), with all but three of the domestic exposures being due to bat-strain rabies.¹⁰ The first case of raccoon-strain rabies in a human occurred in Virginia in 2003. Animal bite surveillance provides the first step in identifying the need for PEP in order to prevent rabies in humans.¹⁸

The incidence of animal bites is considerably higher among children, particularly those five to nine years of age.⁴⁻⁶ Incidence decreases as age increases. Injuries inflicted by dogs are most common (80%-90%), with cats being the next most common species involved.^{3,4} Doberman pinschers, German shepherds, and pit bull terriers are the most common purebred canines implicated in fatal attacks.^{6,11}

B. Disease Surveillance Objectives

1. Determine risk characteristics of patients potentially exposed to animals with rabies.
2. Determine or understand the disposition (euthanized, 10-day quarantine, LTFU) of the biting animal.
3. Assess appropriateness of PEP administration and quality of management of animal bites.

C. Surveillance Indicators

1. Proportion of animal bites reported to the health department in a timely manner.
2. Proportion of cases with complete demographic, exposure information, reported species, and known outcomes for patient and animal.
3. Proportion of animal bite patients that completed a recommended PEP administration.
4. Variables for Performance Evaluation:
 - a. Demographic information
 - i. County of Residence/Zip Code
 - ii. Age
 - iii. Sex
 - iv. Race

Office of Epidemiology and Prevention Services

Division of Communicable Disease Epidemiology

350 Capitol Street Room 125, Charleston, WV 25301-3715

Phone: (304) 558-5358 ext. 2 Fax: (304) 558-6335 www.oeps.wv.gov

Animal Bite/Potential Rabies Exposure Surveillance and Investigation Protocol

- v. Ethnicity
- b. Exposure information
 - i. Date of exposure
 - ii. Type of exposure
 - iii. Bodily location of exposure (for bites)
 - iv. Exposure provoked?
- c. Species information
 - i. Species of the exposing animal
 - ii. Classification of the animals as owned or non-owned (e.g., wild or stray)
 - iii. For owned animals, rabies vaccination status
- d. Known outcomes for victim
 - i. PEP not indicated
 - ii. Patient initiated PEP
 - iii. Patient completed PEP
 - iv. Patient hospitalization
 - v. Patient death
 - vi. Patient previously vaccinated
- e. Known outcomes for exposing animal
 - i. Animal confined
 - ii. Animal outcome
 - iii. Animal tested
 - iv. Animal test results (if tested)

Office of Epidemiology and Prevention Services

Division of Communicable Disease Epidemiology

350 Capitol Street Room 125, Charleston, WV 25301-3715

Phone: (304) 558-5358 ext. 2 Fax: (304) 558-6335 www.oeps.wv.gov

Animal Bite/Potential Rabies Exposure Surveillance and Investigation Protocol

V. REFERENCES

1. Centers for Disease Control and Prevention. Human rabies prevention- United States, 2008. *MMWR Morb Mortal Wkly Rep* 2008 57(Early Release: May 7, 2008):1-30.
2. Centers for Disease Control and Prevention. Update: Raccoon Rabies Epizootic—United States and Canada, 1999. *MMWR Morb Mortal Wkly Rep* 2000 49(02):31-5.
3. Goldstein EJ. Bite wounds and infections. *CID* 1992 14(3):633-8.
4. Garcia, VF. Animal bites and pasteurized infections. *Pediatrics in Review*. 1997 18:127-130.
5. Langley, RL. Human fatalities resulting from dog attacks in the United States, 1979-2005. *Wilderness Environ Med*. 2009 Spring;20(1):19-25.
6. Centers for Disease Control and Prevention. Dog-bite-related fatalities -- United States, 1995-1996. *MMWR Morb Mortal Wkly Rep* 1997 46(21):463-466.
7. Balsamo GA, Ratard R, Claudet A. The epidemiology of animal bite, scratch, and other potential rabies exposures, Louisiana. *J La State Med Soc*. 2009 161(5):260-5.
8. Moore DA, Sicho WM, Hunter A, Miles T. Animal bite epidemiology and surveillance for rabies postexposure prophylaxis. *J Am Vet Med Assoc*. 2000 15;217(2):190-4.
9. Dvorak G, Rovid-Spickler A, Roth JA (editors). 2008. Handbook for zoonotic diseases of companion animals. Center for Food Security and Public Health, Iowa State University, Ames, IA.
10. Centers for Disease Control and Prevention. Rabies surveillance in the U.S.: human rabies. Available at:
http://www.cdc.gov/rabies/location/usa/surveillance/human_rabies.html Accessed 4 March 2022.
11. Sacks JJ, Sinclair L, Gilchrist J. Breeds of dogs involved in fatal human attacks in the United States between 1979 and 1998. *JAVMA* 2000 Vol 217(6):836-840.
12. Beran, GW (editor). 1994. Handbook of zoonoses, 2nd ed. CRC Press, Boca Raton, FL.

Office of Epidemiology and Prevention Services

Division of Communicable Disease Epidemiology

350 Capitol Street Room 125, Charleston, WV 25301-3715

Phone: (304) 558-5358 ext. 2 Fax: (304) 558-6335 www.oeps.wv.gov

Animal Bite/Potential Rabies Exposure Surveillance and Investigation Protocol

13. Heymann, DL (editor). 2008. Control of communicable diseases manual, 19th ed. APHA, Washington DC.
14. Brook L. Management of human and animal bite wound infection: an overview. *Curr Infect Dis Rep.* 2009 11(5):389-95.
15. Freshwater A. Why your housecat's trite little bite could cause you quite a fright: a study of domestic felines on the occurrence and antibiotic susceptibility of *Pasteurella multocida*. *Zoonoses Public Health* 2008 55(8-10):507-13.
16. Blanton JD, Robertson K, Palmer D, Rupprecht CE. Rabies surveillance in the United States during 2008. *JAVMA* 2009 235(6):676-89.
17. Centers for Disease Control and Prevention. Use of a reduced (4-dose) vaccine schedule for postexposure prophylaxis to prevent human rabies: recommendations of the Advisory Committee on Immunization Practices. *MMWR Morb Mortal Wkly Rep* 2010 59(RR-2):1-12.
18. Centers for Disease Control and Prevention. What are the signs and symptoms of rabies? Available at <https://www.cdc.gov/rabies/symptoms/index.html> Accessed 4 March 2022.