

**West Virginia Sexually Transmitted Diseases  
Chlamydia, Gonorrhea and Total Early Syphilis By Age and Gender  
2013 - 2017**

<b>CHLAMYDIA</b>															
	2013			2014			2015			2016			2017		
Age	Male	Female	Total	Male	Female	Total	Male	Female	Total	Male	Female	Total	Male	Female	Total
<10	2	4	6	2	1	3	0	1	1	0	0	0	2	4	6
10-14	2	32	34	2	37	39	4	31	35	2	30	32	3	20	23
15-19	271	1,257	1,528	247	1,097	1,344	281	1,201	1,482	258	1,103	1,361	244	935	1,179
20-24	686	1,589	2,275	629	1,508	2,137	682	1,465	2,147	599	1,381	1,980	555	1,161	1,716
25-29	293	458	751	286	469	755	307	505	812	328	446	774	285	389	674
30-34	141	181	322	104	158	262	122	163	285	132	165	297	124	139	263
35-39	63	78	141	60	74	134	56	83	139	65	74	139	76	55	131
40-44	36	27	63	24	27	51	29	23	52	34	34	68	36	40	76
>44	33	20	53	32	37	69	50	20	70	44	23	67	39	21	60
Unknown	0	0	0	0	1	1	0	0	0	0	0	0	0	0	0
<b>Total</b>	<b>1,527</b>	<b>3,646</b>	<b>5,173</b>	<b>1,386</b>	<b>3,409</b>	<b>4,795</b>	<b>1,531</b>	<b>3,492</b>	<b>5,023</b>	<b>1,462</b>	<b>3,256</b>	<b>4,718</b>	<b>1,364</b>	<b>2,764</b>	<b>4,128</b>
<b>GONORRHEA</b>															
	2013			2014			2015			2016			2017		
Age	Male	Female	Total	Male	Female	Total	Male	Female	Total	Male	Female	Total	Male	Female	Total
<10	0	0	0	1	0	1	0	1	1	0	0	0	0	0	0
10-14	1	3	4	1	7	8	0	2	2	0	2	2	1	3	4
15-19	56	129	185	39	121	160	33	105	138	46	92	138	62	164	226
20-24	173	222	395	132	174	306	102	129	231	140	159	299	184	212	396
25-29	106	97	203	86	90	176	91	72	163	115	83	198	127	120	247
30-34	69	51	120	56	46	102	65	25	90	55	41	96	101	56	157
35-39	41	21	62	22	21	43	37	14	51	44	24	68	76	35	111
40-44	34	14	48	16	9	25	24	6	30	39	12	51	43	21	64
>44	48	11	59	42	9	51	55	7	62	71	18	89	84	12	96
<b>Total</b>	<b>528</b>	<b>548</b>	<b>1,076</b>	<b>395</b>	<b>477</b>	<b>872</b>	<b>407</b>	<b>361</b>	<b>768</b>	<b>510</b>	<b>431</b>	<b>941</b>	<b>678</b>	<b>623</b>	<b>1,301</b>
<b>EARLY SYPHILIS*</b>															
	2013			2014			2015			2016			2017		
Age	Male	Female	Total	Male	Female	Total	Male	Female	Total	Male	Female	Total	Male	Female	Total
<10	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
10-14	0	0	0	0	0	0	0	1	1	0	0	0	0	0	0
15-19	0	0	0	3	2	5	2	3	5	6	2	8	5	8	13
20-24	7	2	9	15	2	17	18	3	21	21	13	34	12	5	17
25-29	5	2	7	11	4	15	16	3	19	9	5	14	17	10	27
30-34	0	1	1	4	3	7	10	5	15	9	4	13	9	4	13
35-39	3	0	3	7	2	9	10	1	11	8	0	8	4	1	5
40-44	1	0	1	1	2	3	5	2	7	10	1	11	3	0	3
>44	4	0	4	6	0	6	13	2	15	12	3	15	16	1	17
<b>Total</b>	<b>20</b>	<b>5</b>	<b>25</b>	<b>47</b>	<b>15</b>	<b>62</b>	<b>74</b>	<b>20</b>	<b>94</b>	<b>75</b>	<b>28</b>	<b>103</b>	<b>66</b>	<b>29</b>	<b>95</b>

\* Early syphilis includes primary, secondary and early latent (less than 1 year) stages of infection.

\*\* Two probable cases of congenital syphilis are not counted in this report: one stillborn and one live birth.