

# Rifabutin

## *Patient information sheet*

### **Why you are taking the medication:**

Rifabutin is used alone or with other drugs to treat tuberculosis and to prevent people from getting sick if they have had contact with tuberculosis bacteria

### **Let your health care provider know right away if you have any of the following issues:**

- Unusual tiredness or loss of appetite
- Severe abdominal upset
- Fever or chills
- Any eye pain, change in vision, or sensitivity to light
- Pain or swelling in the joints
- Yellowing of the skin or eyes or dark urine
- Nausea or Vomiting

### **Instructions for taking medication:**

- Best taken without food; if it bothers your stomach, try taking it with a small amount of food
- It is normal for your urine, tears, and other secretions to turn a brownish- orange color when taking this medicine
- Soft contact lenses may become discolored while you are on this medicine
- Use a second form of birth control to prevent pregnancy while taking this medicine because Rifabutin may decrease the effectiveness of oral hormone-based birth control methods
- As with all medications, it is important to take the full course of the medicine, even if you begin to feel better
- Rifabutin interacts with many types of medications. It is very important to tell your healthcare provider about ANY medication you take including over the counter medication

### **Contact information for your health department:**

---

#### **WV Division of Tuberculosis Elimination**

350 Capitol Street, Room 125, Charleston, WV 25301

304-558-3669(ph) 304-558-1825(fax)

[www.dhhr.wv.gov/oeps/tuberculosis](http://www.dhhr.wv.gov/oeps/tuberculosis)