Proportion of Vaccine-Preventable Disease Cases with Complete Information for Select Indicators, 2016-20222, West Virginia

SURVEILLANCE INDICATOR:		Target	2016	2017	2018	2019	2020	2021	2022
1	The proportion of Haemophilus influenzae invasive disease cases among children under 5 years of age with complete vaccination history.	90%	50% (1/2)	60% (3/5)	100% (3/3)	100% (4/4)	50% (1/2)	100% (3/3)	100 (5/5)
2	The proportion of Haemophilus influenzae isolates from cases under 5 years of age that were serotyped.	90%	100% (2/2)	80% (4/5)	67% (2/3)	75% (3/4)	50% (1/2)	100% (3/3)	80% (4/5)
3	The proportion of measles cases with complete vaccination history.	100%	n/a	n/a	n/a	n/a	n/a	n/a	n/a
4	The proportion of measles cases or chains of transmission that have an imported source.	100%	n/a	n/a	n/a	n/a	n/a	n/a	n/a
5	The proportion of meningococcal cases with complete vaccination history.	90%	75% (3/4)	100% (3/3)	100% (1/1)	NA (0/0)	100% (1/1)	0% (0/1)	100% (1/1)
6	The proportion of meningococcal cases with known serogroup.	90%	100% (4/4)	100% (3/3)	100% (1/1)	NA (0/0)	100% (1/1)	0% (0/1)	100% (1/1)
7	The proportion of mumps cases for which appropriate clinical specimens were obtained and submitted to the laboratory.	80%	80% (4/5)	100% (3/3)	100% (3/3)	100% (2/2)	100% (1/1)	100% (1/1)	100% (2/2)
8	The proportion of mumps cases with complete vaccination history.	90%	100% (5/5)	100% (3/3)	100% (3/3)	100% (2/2)	100% (1/1)	0% (0/1)	100% (2/2)
9	The proportion of pertussis cases from which clinical specimens are obtained.	60%	100% (26/26)	100% (12/12)	100% (25/25)	89% (33/37)	100% (32/32)	100% (4/4)	50% (7/14)
10	The proportion of probable and confirmed pertussis cases meeting the clinical case definition that is laboratory confirmed.	70%	77% (20/26)	58% (7/12)	88% (22/25)	62% (23/37)	38% (12/32)	100% (4/4)	50% (7/14)
11	The proportion of cases confirmed by isolation of B. pertussis by culture.	2%	4% (1/26)	0% (0/12)	8% (2/25)	0% (0/37)	3% (1/32)	0% (0/4)	0% (0/14)
12	The proportion of probable and confirmed pertussis cases with a complete vaccination history.	50%	85% (22/26)	67% (8/12)	100% (25/25)	100% (37/37)	100% (32/32)	0% (0/4)	100% (14/14)
13	The proportion of pneumococcal invasive disease cases among children under 5 years of age with complete vaccination history.	90%	42% (5/12)	85% (11/13)	100% (10/10)	93% (13/14)	67% (4/6)	50% (1/2)	92% (12/13)
14	The proportion of pneumococcal isolates from cases of invasive disease under 5 years of age that are serotyped and tested for antibiotic resistance.	80%	58% (7/12)	85% (11/13)	80% (8/10)	71% (10/14)	67% (4/6)	100% (2/2)	54% (7/13)
15	The proportion of confirmed rubella cases among women of child-bearing age with known pregnancy status.	100%	n/a	n/a	n/a	n/a	n/a	n/a	n/a
16	The proportion of confirmed rubella cases that are laboratory confirmed.	100%	n/a	n/a	n/a	n/a	n/a	n/a	n/a
17	Percentage of varicella cases with complete information for age, vaccination history, and severity of disease.	90%	n/a	n/a	n/a	n/a	n/a	n/a	n/a